



Phymatotrichum Root Rot

Guide A-229

Natalie P. Goldberg
Extension Plant Pathologist

Cooperative Extension Service • College of Agriculture and Home Economics



This publication is scheduled to be updated and reissued 11/10.

Diagnosis at a Glance

Causal agent:	Soil-borne fungus, <i>Phymatotrichopsis omnivorum</i>
Hosts:	Over 2,300 species of broadleafed plants
Symptoms:	Slight yellowing and bronzing of leaves Wilt Reddish lesion on crown Rapid death (leaves remain firmly attached)
Signs:	Fungal threads on roots Spore mats around infected plants
Conditions for disease:	Alkaline soils Soils low in organic matter
Management:	Avoid infested locations Plant non-hosts or resistant species

Phymatotrichum root rot, also known as cotton root rot or Texas root rot, is caused by the soil-borne fungus, *Phymatotrichopsis omnivorum*. Of all diseases known to occur on broadleafed plants, this is one of the most destructive and difficult to control. The pathogen has an incredibly wide host range and surefire survival techniques—two features that make management a nightmare for unfortunate growers faced with the disease. Although thoroughly studied since its discovery in 1888, researchers, farmers, and homeowners still have little ammunition in the fight against the disease.

Fortunately this fungus is limited in distribution and does not readily spread from one location to another. The fungus is restricted geographically to the southwestern United States and northern Mexico in alkaline soils with low organic matter. It also occurs only at elevations below 5,000 feet and is typically found in relatively small areas. In New Mexico, it occurs only in the south-

ern counties, and is most prevalent in agricultural areas along the Rio Grande and Pecos rivers. It spreads slowly from plant to plant when a fungal strand from an infected root grows through the soil to a nearby healthy root. It has no means of air-borne spread.

The fungus has unique biological characteristics that contribute to management difficulties. First, *P. omnivorum* has an extremely wide host range, infecting over 2,300 species of unrelated dicotyledonous (broadleafed) plants. The fungus attacks only mature plants; seedlings are not susceptible to the disease. In New Mexico, this is an important disease of cotton, alfalfa, grapes, fruit trees, and many ornamentals including conifers (figs. 1 A, B). The fungus can also survive on roots of native vegetation, such as mesquite, without causing any disease. Second, the fungus persists almost indefinitely in soil. Much of the fungus is found in the top 2 to 6 feet of soil, however sclerotia (fungal survival structures) have been found over 12 feet deep. Another concern is that isolates of the fungus are non-specific. For example, an isolate infecting cotton is also virulent on fruit trees and ornamentals and vice versa. Thus, housing developments in old cotton and alfalfa fields with a history of the disease can be disastrous for urban landscapers.

The fungus is only active in summer months when air and soil temperatures are high. The greatest incidence of disease occurs when soil temperature at 1 foot deep is greater than 80°F and the air temperature in the plant canopy is above 104°F. When environmental conditions are favorable for fungal activity, the pathogen invades plants through their root systems. Infected roots rot and cannot transport water to the above-ground portion of the plant. Symptoms on above-ground plant parts resemble water stress. The first evidence of disease is slight yellowing of the leaves. Leaves quickly turn to a bronze color and begin to wilt. Permanent wilting of the branches can occur very rapidly, as little as two weeks from the first expression of disease. The tree dies with leaves remaining firmly attached (fig. 2). In some cases, the tree wilts so quickly that the leaves hardly change color, though they will become dry and brittle. A red-




Figure 1A. *Phymatotrichum* root rot in alfalfa



Figure 1B. *P. omnivorum* in peaches



Figure 2. Plant killed by *P. omnivorum*; note leaves still attached



Figure 3. Fungal strands on infected roots

dish lesion around the crown of the plant may develop on trees killed by the fungus.

Evidence of *P. omnivorum* can also be found on or near infected trees. The pathogen produces fungal strands on the surface of infected roots (fig. 3). These strands look like threads and are visible with a good hand lens. When strands are observed under a compound microscope, cruciform (cross-shaped) hyphae unique to this fungus can be seen (fig. 4). Another sign is the formation of a white- to tan-colored spore mat on the surface of the soil around infected plants (fig. 5). Spore mats develop during periods of high moisture. The spores in these mats have never been germinated and are considered to have no function in survival or

infection of the pathogen. Therefore, spore mats do not spread disease, but are evidence of its presence.

Research to control this disease has been extensive, yet there are no good control methods. There is no resistance or tolerance to this disease in most of the commonly infected hosts. The best recommendation is to avoid land known to be infested with the fungus. Furthermore, fruit and nut orchards should not be planted in old cotton or alfalfa fields. If mesquite land is to be cleared for orchard planting, it may be worth the time and money to replant the area in cotton. Plants with a tap root like this are infected and show symptoms in one season, while trees with fibrous root systems do not show disease for some time after planting. If no symp-



Figure 4. Characteristic cruciform hyphae



Figure 5. Spore mat

toms develop on this test crops, growers can be relatively sure the land is safe for growing ornamentals or fruit trees.

The ability of the fungus to survive deep in soil has eliminated the possibility of using fungicides and fumigants to control the disease because these materials can only penetrate a limited distance into the soil. Fumigating infested planting holes in areas where trees have died will usually only delay the onset of disease in newly planted trees. When a replanted tree's roots reach deep enough in the soil, they will contact the surviving fungus and succumb to the disease.

There are practices that can be used to try to alter the soil environment so it no longer favors *P. omnivorum*. These practices may help reduce the effect of the pathogen if they are followed every year. If not, the soil environment returns to its typical state (high pH and low organic matter) and again favors the fungus. The procedure consists of loosening the soil in a broad and comparatively shallow basin just beyond the drip line around infected trees. The area is then covered to a depth of 2 inches with manure or similar organic matter. Layered on top of the organic matter is ammonium sulfate and sulfur, each at a rate of 1 lb/10 sq ft. The basin should be immediately flooded with enough water to wet the soil to a depth of 3 feet. This high level of soil moisture must be maintained for several weeks. If trees are treated before permanent wilting, they may recover. Known root rot-infested areas should be treated every year in March or April.

Some success reducing the effect of the disease has been achieved by growing and incorporating a green manure cover crop over the orchard floor. This helps to stimulate vigorous rooting of the trees, enabling them

to better withstand disease pressure. Additionally, the incorporation of the cover crop into the soil may help to stimulate soil microflora, which compete with *P. omnivorum*.

Finally, chemical barriers have been used to prevent the spread of disease from one tree to another. Using sulfur in trenches 4 to 6 inches wide and 4 to 6 feet deep around the outside of the drip line of infected trees has prevented the spread of *P. omnivorum* for over 7 years.

In areas known to be infested with *P. omnivorum*, the only sure control is to plant non-hosts. Unfortunately, the extensive host range of the pathogen limits choices. Monocots, such as corn, small grains, and all grasses are immune. Dicots grown as annuals often escape the disease. Additionally, most native desert plants are apparently tolerant to the disease.

The best thing that can be said about this fungus is that it is not widespread. It is found only in small pockets and does not spread much from its point of origin. There are no viable spores to help spread the disease from one location to another. The only known spread between plants occurs when fungal threads from infected roots contact healthy roots.

New Mexico State University is an equal opportunity/affirmative action employer and educator. NMSU and the U.S. Department of Agriculture cooperating.